

# Clustering features of the ${}^7\text{Be}$ nucleus in relativistic fragmentation

N. K. Kornegrutsa, D. A. Artemenkov, P. I. Zarubin,\* and I. G. Zarubina

*Joint Insitute for Nuclear Research, Dubna, Russia*

R. R. Kattabekov

*Institute for Physics and Technology,*

*Uzbek Academy of Sciences, Tashkent, Republic of Uzbekistan*

K. Z. Mamatkulov

*Djizak State Pedagogical Institute, Djizak, Republic of Uzbekistan*

(Dated: October 21, 2014)

## Abstract

Charge topology of fragmentation of 1.2 A GeV  ${}^7\text{Be}$  nuclei in nuclear track emulsion is presented. The dissociation channels  ${}^4\text{He} + {}^3\text{He}$ ,  $2{}^3\text{He} + \text{n}$ ,  ${}^4\text{He} + 2{}^1\text{H}$  are considered in detail. It is established that the events  ${}^6\text{Be} + \text{n}$  amount about to 27 % in the channel  ${}^4\text{He} + 2{}^1\text{H}$ .

PACS numbers: 21.45.+v, 23.60+e, 25.10.+s

---

\*Electronic address: zarubin@lhe.jinr.ru; URL: <http://becquerel.jinr.ru>

## I.

Stacks of pellicles of nuclear track emulsion provide a special opportunity to explore clustering of light nuclei (reviewed in [1]). The presented results on dissociation of  ${}^7\text{Be}$  nuclei are demonstrate the progress in research carried out by the BECQUEREL Collaboration. The  ${}^7\text{Be}$  nucleus is a source for the study of the states  ${}^3\text{He} + {}^4\text{He}$ ,  ${}^3\text{He} + {}^3\text{He} + n$ ,  ${}^6\text{Li} + p$  and  ${}^6\text{Be} + n$ . The pattern of fragmentation is important for understanding of the structure features of the nuclei  ${}^8\text{B}$ ,  ${}^9\text{C}$  and  ${}^{12}\text{N}$  because the  ${}^7\text{Be}$  nucleus plays the role of a core in them.

Nuclear track emulsion was irradiated at the Nuclotron of the Joint Institute for Nuclear Research (JINR, Dubna) by a mixed beam of  ${}^7\text{Be}$ ,  ${}^{10}\text{C}$ , and  ${}^{12}\text{N}$  nuclei which was created by selecting products of charge-exchange and fragmentation processes involving  ${}^{12}\text{C}$  nuclei accelerated to an energy of 1.2 GeV per nucleon [2], [3], [4], [5]. Viewing of the exposed pellicles and the track classification made it possible to establish the charge topology of the  ${}^7\text{Be}$  nucleus. Peripheral fragmentation distribution of the 289 found events  $N_{ws}$  not accompanied by target fragments (“white ”stars) is presented in Table 1 over the fragmentation channels as well as 380 events  $N_{tf}$  accompanied by target fragments.

The distribution of the 79 events  $2\text{He}$  which were successfully identified by multiple scattering is presented in Table 2. It gives an idea about the relationship configurations  ${}^3\text{He} + {}^4\text{He}$  and  $2{}^3\text{He} + n$  in the  ${}^7\text{Be}$  structure, as the identification was carried out without bias. The channel  ${}^3\text{He} + {}^4\text{He}$  dominates over  $2{}^3\text{He}$  indicating on a higher probability of the two-body configuration  ${}^3\text{He} + {}^4\text{He}$  in the  ${}^7\text{Be}$  structure compared to  $2{}^3\text{He} + n$ . The probability of the  $2{}^3\text{He} + n$  channel is significant, amounting to about 30%.

The distribution of the events  $2{}^3\text{He}$  and  ${}^3\text{He} + {}^4\text{He}$  over the excitation energy  $Q_{2He}$  defined as a difference of the invariant mass of the fragmenting system and the sum of the fragment masses is shown in Fig. 1. The values  $Q_{2H}$  of the channel  ${}^3\text{He} + {}^4\text{He}$  are distributed in the range covering the known levels of the  ${}^7\text{Be}$  nucleus excitation.

One of the tasks of this study consisted in searching for narrow pairs  $2{}^3\text{He}$  with values  $Q_{2{}^3He}$  in a range of 100 – 200 keV the indication to which was obtained for dissociation  ${}^9\text{C} \rightarrow 2{}^3\text{He}$ . The obtained distribution includes four events with values in the range of 200–400 keV (Fig. 1, dotted histogram in insertion). These data do not exclude a possible existence of the resonant state  $2{}^3\text{He}$  discussed in [6].

TABLE I: Distribution over the dissociation channels of  ${}^7\text{Be}$  nuclei for “white” stars  $N_{ws}$  and events with target fragments or produced mesons  $N_{tf}$ .

	Channel $2\text{He} \text{ He} + 2\text{H} \text{ 4H Li} + \text{H}$			
$N_{ws}$	115	157	14	3
$N_{tf}$	154	226	-	-

TABLE II: Distribution over the dissociation channels of  ${}^7\text{Be}$  nuclei for “white” stars  $N_{ws}$  and events with target fragments or produced mesons  $N_{tf}$ .

	Channel ${}^3\text{He} + {}^4\text{He} \text{ } {}^3\text{He} + {}^3\text{He}$	
$N_{ws}$	32	14
$N_{tf}$	24	9

There is an opportunity of  ${}^7\text{Be}$  fragmentation via an unstable  ${}^6\text{Be}$  nucleus with a threshold 1.37 MeV above  ${}^4\text{He} + 2\text{p}$ . Fig. 2 shows distribution of events  ${}^4\text{He} + 2\text{p}$  over the difference of the invariant mass of the produced  $\alpha$ -particle and two protons and their mass sum  $Q_{{}^4\text{He}+2\text{p}}$ . The region  $Q_{{}^4\text{He}+2\text{p}} < 6$  MeV indicates on the presence of about 27% events  ${}^7\text{Be} \rightarrow {}^6\text{Be} \rightarrow {}^4\text{He} + 2\text{p}$ . Thus, contribution of the configuration  ${}^6\text{Be} + \text{n}$  to the  ${}^7\text{Be}$  structure is estimated at a level of  $8 \pm 1$  %.

The question about the contribution of the  ${}^5\text{Li}$  resonance decaying to  $\alpha + \text{p}$  with an energy of 1.69 MeV and width of 1.5 MeV has a significance independent of  ${}^6\text{Be}$  since the production threshold of  ${}^5\text{Li} + \text{p}$  is 0.35 MeV higher than the one of the ground state  ${}^6\text{Be}$ . Despite of the absence of a clear signal the distribution  $Q_{\alpha\text{p}}$  (Fig. 3) does not contradict to a possible contribution of  ${}^5\text{Li}$  decays.

This work was supported in part by the Russian Foundation for Basic Research (project no. 12-02-00067) and by grants from the plenipotentiaries of Bulgaria and Romania at the Joint Institute for Nuclear Research (Dubna).




FIG. 1: Distribution of events  ${}^7\text{Be} \rightarrow {}^3\text{He} + {}^4\text{He}$  and  ${}^{23}\text{He}$  over the excitation energy  $Q_{2He}$  (solid and dotted histograms, respectively). Histograms for values  $Q_{2He} < 1$  MeV are on the insertion and  ${}^{23}\text{He}$  over the excitation energy  $Q_{2He}$  (solid and dotted histograms, respectively). Histograms for values  $Q_{2He} < 1$  MeV are on the insertion.




FIG. 2: Distribution of events  ${}^7\text{Be} \rightarrow {}^4\text{He} + 2p$  over the excitation energy  $Q_{4He+2p}$ .




FIG. 3: Distribution of events  ${}^7\text{Be} \rightarrow {}^4\text{He} + 2p$  over the excitation energy  $Q_{\alpha p}$  (events related to the  ${}^6\text{Be}$  decays are excluded from this histogram.)

- 
- [1] Zarubin P. I., Lect. Notes in Phys., 875, Clusters in Nuclei, Vol. 3. Springer International Publishing, pp 51-93. 2013; arXiv:1309.4881.
  - [2] Rukoyatkin P. A. et al., EPJ ST 162, 267 (2008); arXiv:1210.1540.
  - [3] Kattabekov R. R. et al., Phys. Atom. Nucl. 73, 2110 (2010); arXiv:1104.5320.
  - [4] Mamatkulov K. Z. et al. Phys. At. Nucl., 2013, Vol. 76, No. 10, pp. 1224-1229; arXiv:1309.4241.
  - [5] Kattabekov R. R. et al. Phys. At. Nucl., 2013, Vol. 76, No. 10, pp. 1219-1223; arXiv:1310.2080.
  - [6] Artemenkov D. A. et al., J. Phys. Conf. Ser. 337 (2012) 012019 arXiv:1105.3813.